

Smectic Order in Double-Twist Cylinders

Randall D. Kamien[†]

*Department of Physics and Astronomy, University of Pennsylvania,
Philadelphia, PA 19104, USA*

I propose a double-twist texture with local smectic order, which may have been seen in recent experiments [1]. As in the Renn-Lubensky TGB phase, the smectic order is broken only through a lattice of screw dislocations. A melted lattice of screw dislocations can produce a double-twist texture as can an unmelted lattice. In the latter case I show that geometry only allows for certain angles between smectic regions. I discuss the possibility of connecting these double-twist tubes together to form a smectic blue phase.

4 November 1996; revised 13 February 1997

[†] kamien@dept.physics.upenn.edu

1. Introduction

Since the prediction [2] and discovery [3] of the defect-laden, Renn-Lubensky, twist-grain-boundary ground state of a chiral smectic, it has become *de rigueur* to consider defected ground states in liquid crystals [4,5]. While it is possible to introduce defects into smectics or columnar phases to produce planar chiral textures like the cholesteric or twisted hexatic [6], there has been no attempt to construct the more complicated blue phase textures [7] in a similar fashion. Recent experiments [1], however, have suggested that there exist blue phases with some sort of smectic order. In this letter, I propose a general defect structure that can produce smectic double-twist. The smectic order may be only *short-ranged* as is seen in the N_L^* phase of some liquid crystals [8,9]. I will first discuss this case, which is likely to be the generic case, and will then consider the possibility of having long-range smectic order. In this case, the defects will form a regular lattice. It will be shown that because of the periodicity imposed by the cylindrical structure of a double-twist tube the allowed rotation angles fall into a discrete class similar to, but not as restrictive as, moiré angles [4].

I start by recalling the general structure of a twist-grain-boundary in a smectic liquid crystal. The free energy for a smectic- A can be written to quadratic order in terms of the layer displacement variable $u(\mathbf{x})$ and the nematic director fluctuation $\delta\mathbf{n}(\mathbf{x}) = \mathbf{n} - \hat{\mathbf{z}}$ [10]:

$$F = \frac{1}{2} \int d^3x B(\partial_z u)^2 + B'(\nabla_\perp u - \delta\mathbf{n})^2 + K_1(\nabla_\perp \cdot \delta\mathbf{n})^2 + K_2(\nabla_\perp \times \delta\mathbf{n})^2 + K_3(\partial_z \delta\mathbf{n})^2, \quad (1.1)$$

where B and B' are bulk moduli and K_i are the Frank elastic constants. In the ground state, $\nabla_\perp u = \delta\mathbf{n}$, and so $\nabla_\perp \times \delta\mathbf{n} \equiv 0$. Thus a smectic cannot have any twist. However, as pointed out by Renn and Lubensky, the smectic can twist if u is not single-valued: in other words if there are screw dislocations in the smectic structure [2]. Each screw dislocation has a finite energy per unit length and so there will be a balance between the cost of a defect and the amount of twist produced by the structure. Above some critical

chirality the defects will proliferate and organize themselves into a sequence of twist-grain-boundaries. In the twist-grain-boundary (TGB) state of smectic-*A* liquid crystals, the nematic director rotates around a pitch axis $\hat{\mathbf{P}}$ which lies in the plane of the smectic layers. The defected structure consists of regions of perfect smectic layers, infinite in the two directions perpendicular to $\hat{\mathbf{P}}$ and of extent d' along $\hat{\mathbf{P}}$. Between each successive slab there is a twist-grain-boundary consisting of parallel screw dislocations with a uniform spacing d . Each grain boundary effects a twist of the director by $\Delta\theta = 2\sin^{-1}(b/2d)$ [11] where b is the Burgers vector which must be an integral multiple of the layer spacing a . The lowest energy grain boundary for a given rotation angle will consist of screw dislocations of unit strength, *i.e.* $b = a$. In the end, the TGB state has the director structure of a cholesteric (with discretized jumps in angle) while at the same time having long-range smectic order.

As temperature increases, if the smectic order does not melt first, the lattice of screw dislocations can melt, in analogy with the melted flux-line liquid of superconductors [12]. This N_L^* phase [8] is thermodynamically the same as the cholesteric phase and has a uniformly twisting director structure and is composed of a melt of dislocation lines which themselves follow the director, as if they were polymers in a polymer cholesteric phase [9]. While there is no long-range smectic order, there is short range smectic order which can be seen via X-ray scattering and through calorimetry: before the TGB state appears there is an absorption of heat which is required to build up the local smectic order.

How can either of these dislocated structures be deformed to produce double-twist? Since the smectic order is locally preserved, there must be a defect structure which produces an average double-twist, though possibly via non-uniform jumps. Such a structure is shown in Figures 1 and 2, in two cross-sectional views. In this structure the layer normal has discretized double-twist as does the tangent vector field of the screw dislocations. The structure of the double-twist tube is identical to the usual TGB structure when *wrapped* around a central smectic region. While the grain boundary structure is highly

distorted near the center of the tube, it is clear that at larger radii the local structure at a grain boundary is identical to the structure in the more usual cholesteric-like TGB state. Thinking of the grain-boundaries as smectic layers, the grain boundaries adopt a standard jelly-roll texture [13]. In the next section I will discuss some of the geometric details required to build such a structure. Section 3 will discuss the energetics of the double-twist tube. Finally, in section 4, I will consider the possibility of a smectic blue phase.

2. Geometric Details: Layer Spacing and Lock-In Angles

While the above geometric picture of the defect structure captures the essence of the construction, the details are significantly more complex. One concern arises because the smectic layer tilt and spacing naturally change as a function of radius. In one undefected smectic region the spacing between the layers along the tube axis (*not* perpendicular to the layers) must be a constant δ as shown in Figure 3. Thus if $\theta(R)$ is the angle that the layer normal makes with the tube axis then $\delta = R \tan \theta(R)$ so that R and $\theta(R)$ obey

$$\frac{\tan \theta(R_1)}{\tan \theta(R_2)} = \frac{R_2}{R_1}. \quad (2.1)$$

Similarly, the layer spacing is related to $\theta(R)$ by

$$\frac{a(R_1)}{a(R_2)} = \sqrt{\frac{1 + \tan^2 \theta(R_2)}{1 + \tan^2 \theta(R_1)}} \equiv \frac{\cos \theta(R_1)}{\cos \theta(R_2)}, \quad (2.2)$$

since $\delta \cos \theta(R) = a(R)$. As $R_2 \rightarrow \infty$ for fixed R_1 , $\tan \theta(R_2) \rightarrow 0$ and $a(R_2) \rightarrow a$, the equilibrium layer spacing (as it always is for even *single* screw dislocations). Both (2.1) and (2.2) come from the equation of the helicoidal surface. In one sweep around the azimuthal direction, the surface, given in cylindrical coordinates by the height function $h(r, \phi) = \alpha \phi$, rises by $2\pi\alpha$, *independent* of the radius. This means that the layer normal changes its angle according to (2.1) and the layer spacing according to (2.2).

The second issue is the periodicity of the structure because it is a cylinder: at radius R the structure must be $2\pi R$ periodic. In particular, when wrapping smectic layers and

grain boundaries around a double-twist cylinder, one must constrain the number of layers or defects to be integral. At a given radius R , assume that the smectic layers have rotated by θ_0 from the center. Additionally, at this same radius add a grain boundary that rotates the layers by $\Delta\theta$ with a dislocation spacing $d(R)$ and with all defects having Burgers vector $b = a(R)$, where $a(R)$ is the layer spacing. If there are n smectic layers in the subsequent smectic region and m defects then trigonometry leads to:

$$\frac{a(R)}{2d(R)} = \sin\left(\frac{\Delta\theta}{2}\right) \quad (2.3a)$$

$$\frac{md(R)}{2\pi R} = \cos\left(\theta_0 + \frac{\Delta\theta}{2}\right) \quad (2.3b)$$

$$\frac{na(R)}{2\pi R} = \sin(\theta_0 + \Delta\theta) \quad (2.3c)$$

where the angle $\theta_0 + \Delta\theta/2$ in (2.3b) arises because the screw dislocations lie symmetrically between the two abutting smectic regions. Eliminating R , $a(R)$ and $d(R)$ from (2.3) leads to:

$$\frac{m}{n} = 1 - \frac{1}{\cos \Delta\theta + \cot \theta_0 \sin(\Delta\theta)} \quad (2.4)$$

The left-hand side of (2.4) is a rational number m/n . While there exists an entire discrete class of solutions to this Diophantine equation, one class of solutions has $\cos \Delta\theta = p_1/q_1$ and $\cot \theta_0 = p_2/(q_2\sqrt{q_1^2 - p_1^2})$ for p_1, p_2, q_1, q_2 integers. Note that if the centermost region contains a single screw dislocation down the center of the double-twist tube then, in order to satisfy (2.4), the first twist-grain-boundary will occur at some radius R such that $\tan \theta_c = (a/2\pi R)$ is of the form $q_2\sqrt{q_1^2 - p_1^2}/p_2$.

Having a class of angles I may now construct many double-twist cylinders that have the layer normal at the center parallel to the cylinder and the layer normal at the edge of the cylinder rotated by $\pi/4$ from this direction. These are the basic constituents of the blue phases. At each grain boundary θ_0 in (2.3b, c) and (2.4) is the sum of the center angle of rotation θ_c and the sum of all the consecutive rotation angles, so if the i th boundary is

at a radius $R = \rho_i$, then $\theta_{0,i} = \theta_c + \sum_{j=1}^{i-1} \tan^{-1}[(\rho_j/\rho_{j+1}) \tan(\Delta\theta_j)]$. Taking into account the natural straightening of the layer normals (2.1) I can choose radii and angles so that at some final radius ρ the layer normal can have the desired direction. Because the tangent angle as a function of radius is also fixed by the geometry of the screw dislocation (*i.e.*, $\tan[\theta_c(R)] = b/[2\pi R]$) (2.3) may be more difficult, if not impossible, to satisfy if there is a screw dislocation running up the center of the double-twist tube.

As a simple illustration, let me construct a double-twist tube with layer normal $\pi/4$ at some ultimate radius ρ via two grain boundaries both with a large rotation angle, $\Delta\theta = \sin^{-1}(3/5)$. This corresponds to an even simpler class of angles in which both sine and cosine are rational – precisely the construction that leads to moiré angles. With this choice of $\Delta\theta$, I must now choose $\cot\theta_{0,i}$ to be rational as well. At the first boundary (at radius ρ_1), $\theta_{0,1} = 0$, so from (2.4) the ratio of the number of layers to the number of screw dislocations is $n_1/m_1 = 1$, while (2.3a) gives $a(\rho_1) = d(\rho_1)/\sqrt{10}$. At radius $\rho_2 = 2\rho_1$, (2.1) gives $\tan\theta_{0,2} = \frac{1}{2} \tan(\Delta\theta) = 3/8$ and is rational, as required. Using the same angle $\Delta\theta$ across the grain boundary now gives $n_2/m_2 = 12/7$ and a total tangent of $\tan(\theta_{0,2} + \Delta\theta) = 36/23$. Finally, then, at a radius $\rho = \frac{36}{23}\rho_2 = \frac{72}{23}\rho_1$ the layer normal is at an angle $\pi/4$ from the tube radius. While this example is rather extreme, involving large angles grain boundaries and large radii, it is obvious that more realistic examples could be constructed.

3. Energetics

Let me now consider the energetics of a smectic double-twist tube. In non-layered chiral liquid crystals, double-twist structures are favored over standard planar cholesteric structures due to a trade-off between the saddle splay term [7]

$$F_{ss} = \int d^3x K_{24} \nabla \cdot [(\hat{\mathbf{n}} \cdot \nabla) \hat{\mathbf{n}} - \hat{\mathbf{n}} (\nabla \cdot \hat{\mathbf{n}})], \quad (3.1)$$

and the twist term

$$F_{\text{tw}} = \int d^3x K_2 [\hat{\mathbf{n}} \cdot (\nabla \times \hat{\mathbf{n}}) - q_0]^2. \quad (3.2)$$

In cylindrical coördinates, the double-twist texture $\hat{\mathbf{n}} = \hat{\mathbf{z}} \cos(q_0 \rho/2) + \hat{\phi} \sin(q_0 \rho/2)$. This director configuration has non-zero saddle splay. This would be irrelevant if there were no director defects in the blue phase: F_{ss} is the integral of a total derivative. Thus saddle splay is gained at the surfaces of the cores of nematic defects. Since the value of the saddle splay $2[\partial_x n_y \partial_y n_x - \partial_x n_x \partial_y n_y] = -q_0 \sin(q_0 R)/(2R)$ is non-zero it can lower the energy compared to the planar texture if $K_{24} > 0$. However, there is an energy cost which comes from the fact that $\hat{\mathbf{n}} \cdot (\nabla \times \hat{\mathbf{n}}) = q_0/2 + \sin(q_0 R)/(2R)$ and thus the twist term only vanishes near the center of the double-twist tube, whereas it vanishes everywhere in the planar cholesteric texture. Note that in one double-twist tube that rotates from 0 to $\pi/4$, the saddle-splay per unit length is $F_{\text{ss}}/L = -\pi q_0^2(\sqrt{2} - 1)/\sqrt{2}$.

The same comparison can be made for layered chiral systems. First the energy of the grain boundaries in the double-twist structure must be compared to the same energy in the standard TGB phase. Neglecting interactions between screw dislocations, the energies should be comparable. A detailed calculation of the interaction energy would be interesting but, because of the complex three-dimensional geometry of the dislocations, difficult (it should be noted that this calculation has not yet been done even for the standard TGB state). In some sense this problem is identical to the problem of deciding between the standard cholesteric phase and the blue phase. The standard TGB phase has a cholesteric-like organization of the screw dislocations while the phase proposed here has a blue phase organization of the screw dislocations. The deciding factor between a cholesteric and a blue phase is the presence of an energetic term which favors or disfavors saddle-splay. Thus, it is natural to ask whether topological effects play any additional role in the smectic double-twist texture. In the layered system the nematic disclinations stay the same but now there are a plethora of screw dislocations. At each screw dislocation the layers can pick up surface energies as well. In fact, these surface energies are completely analogous to

the saddle splay: the Gaussian curvature will contribute to the energy at the edges which separate the successive, annular smectic regions. For a surface given by a height function $h(x, y)$, the integrated Gaussian curvature K is:

$$\int \sqrt{g} d^2\sigma K = \int dx dy \frac{\partial_x^2 h \partial_y^2 h - (\partial_x \partial_y h)^2}{[1 + (\nabla_\perp h)^2]^{3/2}}, \quad (3.3)$$

where $\sqrt{g} dx dy$ is the area element of the surface. If the helicoidal surface of each smectic layer in region i is given by the multi-valued height function $h_i(x, y) = \alpha_i \tan^{-1}(y/x)$, then for the region between ρ_i and ρ_{i+1}

$$\int \sqrt{g_i} d^2x K_i = -2\pi N_i [\cos \theta_{0,i+1} - \cos(\theta_{0,i} + \Delta\theta_i)], \quad (3.4)$$

where N_i is the number of helical turns of the i th annular region. If I consider a segment of the double-twist tube of length L , then $N_i a(\rho_{i+1}) = L \cos \theta_{0,i+1}$ and so the total Gaussian curvature per unit length is

$$\frac{F_{\text{Gauss}}}{L} = -2\pi \sum_{i=1}^N \frac{\cos^2 \theta_{0,i+1}}{a(\rho_{i+1})} \left\{ 1 - \frac{a^2(\rho_i)}{a^2(\rho_{i+1})} \right\}. \quad (3.5)$$

Thus the total Gaussian curvature is always negative (which is no surprise: all helicoidal surfaces have negative Gaussian curvature). The smectic free energy will increase or decrease depending on whether the material in question favors or disfavors positive Gaussian curvature, respectively. Obviously the energetics will depend on the particular choice of angles $\Delta\theta_i$ and radii ρ_i . Note that this extra geometric energy of the double-twisted smectic is analogous to the saddle-splay energy of standard blue phases. The energy gain from the surface Gaussian curvature must offset the energy cost of the extra distortions of the smectic layers and the cholesteric order for double-twist to be favored. If the net energy change is negative the smectic will prefer to have double-twist regions instead of the standard TGB state.

4. Space Filling Structures: Blue Phases

The ability to connect the double-twist tubes into one of the standard blue phases depends on whether the smectic order is short- or long-ranged. In the first case, since the N_L^* phase is identical to a cholesteric, there is no equilibrium impediment to constructing the usual blue phase textures. Since there is only short range smectic order the integrity of the smectic layers need not be globally preserved and there can be any number of edges and screws filling the space between the traditional double-twist tubes. Of course, same-sign defects repel each other and so there may be dynamical reasons that a complete blue phase could not form. It is likely that in the work of [1], that this melted defect lattice is what is seen. If a material has a strong tendency to have smectic order then it is reasonable to believe that short-range smectic order appears even in the normal blue phases.

The case of true, long-range smectic order is problematic. Here we will only consider the O^5 blue phase structure though this work could be extended to the other known arrangements of double-twist tubes [7]. The O^5 blue phase has the simple “log-cabin” arrangement of double-twist tubes as shown in Figure 4. If I arrange for a layer on cylinder A to meet a layer on cylinder B at the intersection point Γ then, since the layer normal is at $\pi/4$ with respect to the cylinder axis, that same layer will shift by $(2\pi)\rho/4$ towards Δ . Since Δ is 2ρ further along axis A from Γ , it would be impossible for the meeting layer at Γ to meet the layer at Δ . In Figure 5 I have unwrapped a double-twist tube at the radius ρ . One can see that if the layer spacing is $a = a(\rho)$ then for each integer j there is a layer at $ja(\rho)\sqrt{2} + \pi\rho/2$ along the (dashed) line on which Δ lies. Thus a layer in cylinder A can meet a layer in cylinder C at Δ if $ja(\rho)\sqrt{2} = (2 - \pi/2)\rho$. At the last grain boundary, however, $na(\rho_N) = \sin(\theta_{0,N} + \Delta\theta_N)2\pi\rho_N$. Using (2.1) and (2.2) I find $j/n = (4 - \pi)/(4\pi) \approx 1.4165$. Since j/n is rational, however, the kissing conditions can never be satisfied. However, it would still be possible, though energetically unfavorable, to change the layer spacing from its preferred value near the cylinder surface. In this case some ratio of integers could be found that makes the layer spacing mismatch small.

Even if this deformation occurred the region between the double-twist tubes would have a complicated complexion of screw and edge dislocations in order to match up *all* the smectic layers.

With some experimental input, it may be possible to show that the constraints presented in this letter could almost be satisfied for an observed blue phase with smectic order. It is most likely that if a blue phase texture with smectic order persists it would only have short-range smectic order. The absence of long-range smectic order would facilitate a number of the obstacles to the construction of the double-twist tubes. For instance, the regions between the double-twist tubes could melt and thus alleviate the rational/irrational mismatch. However, the special lock-in angles would still correspond to low energy grain boundaries since at these angles the smectic order is not frustrated by the double-twist texture of the layer normal. I would thus expect that the lock-in angles would be present in any blue phase with long- or short-ranged smectic order.

5. Acknowledgments

It is a pleasure to acknowledge stimulating discussions with T.C. Lubensky, B. Pansu, T. Powers, J. Sethna, H. Stark and an anonymous referee. I wish to thank the Aspen Center for Theoretical Physics, where some of this work was done. This work was supported by NSF Grant DMR94-23114.

References

- [1] B. Pansu, M.L. Hui and H.T. Nguyen, *to appear in* J. Phys. II France (1997).
- [2] S.R. Renn and T.C. Lubensky, Phys. Rev. A **38** (1988) 2132; **41** (1990) 4392.
- [3] J. Goodby, M.A. Waugh, S.M. Stein. R. Pindak, and J.S. Patel, Nature **337** (1988) 449; J. Am. Chem. Soc. **111** (1989) 8119; G. Strajer, R. Pindak, M.A. Waugh, J.W. Goodby, and J.S. Patel, Phys. Rev. Lett. **64** (1990) 13; K.J. Ihn, J.A.N. Zasadzinski, R. Pindak, A.J. Slaney, and J. Goodby, Science **258** (1992) 275.
- [4] R.D. Kamien and D.R. Nelson, Phys. Rev. Lett. **74** (1995) 2499; Phys. Rev. E **53** (1996) 650.
- [5] G. Yan and T.C. Lubensky, University of Pennsylvania Preprint (1996) [cond-mat/9611066].
- [6] R.D. Kamien, J. Phys. II France **6** (1996) 461.
- [7] S. Meiboom, J.P. Sethna, P.W. Anderson and W.F. Brinkman, Phys. Rev. Lett. **46** (1981) 1216; D.C. Wright and N.D. Mermin, Rev. Mod. Phys. **61** (1989) 385.
- [8] T. Chan, C.W. Garland and H.T. Nguyen, Phys. Rev. E **52** (1995) 5000; L. Navailles, C.W. Garland and H.T. Nguyen, J. Phys. II France **6** (1996) 1243.
- [9] R.D. Kamien and T.C. Lubensky, J. Phys. I France **3** (1993) 2131.
- [10] P.G. de Gennes, Solid State Commun. **14** (1973) 997.
- [11] I thank T. Lubensky for discussions on this point. If you cannot arrange an audience with him, see, for instance, P.M. Chaikin and T.C. Lubensky, *Principles of Condensed Matter Physics*, (Cambridge University Press, Cambridge, 1995), Chapter 9.
- [12] D.R. Nelson, Phys. Rev. Lett. **60** (1988) 1973; D.R. Nelson and H.S. Seung, Phys. Rev. B **39** (1989) 9153.
- [13] P.G. de Gennes and J. Prost, *The Physics of Liquid Crystals*, Second Edition, (Oxford University Press, New York, 1993).

Figure Captions

- Fig. 1. Double-twisted smectic structure. The dark lines are the screw dislocations. Note that the layer normal *and* the screw dislocation tangent vectors have a double-twist texture. While the rotation angles, defect density and layer spacing are related, this figure does not take that into account – it only serves as a simple model.
- Fig. 2. A cut-away view of the double-twisted structure of Figure 1 through the center of the tube.
- Fig. 3. Geometrical construction used to derive (2.1) and (2.2). Coming out radially from the center the spacing of the layers along the tube direction must be a constant δ in one smectic slab. Thus $R \tan \theta(R) = \delta$ and $\delta \cos \theta(R) = a(R)$ from which follow (2.1) and (2.2), respectively.
- Fig. 4. The O^5 blue phase structure. The double-twist cylinders are of radius ρ and each side of the cube shown has length 4ρ . The three cylinders that are cut by the visible faces are labeled A , B and C , while two of the kissing points are labeled Γ and Δ .
- Fig. 5. An unwrapped double-twist cylinder. The plane corresponds to the *inside* surface of the cylinder A in Figure 4. The dashed line is that line parallel to the double-twist axis that is at an angle $\pi/2$ from the similar line in which Γ lies. At Γ the cylinder kisses cylinder B , while at Δ it kisses cylinder C .




Figure 1




Figure 2




Figure 3




Figure 4




Figure 5